

PMG Laboratory Spokane

Test Directory

			1
LAB	CODE	TEST NAME	CPT4 CODE
PMGLAB	LAB560	QUAD SCREEN MATERNAL, SERUM Place Of Service:LABCORP RTP	
PMGLAB	LAB2399	QUANTIFERON GOLD Container:Dark green lithium heparin tube Temperature:Refrigerate Place Of Service:PMG-PCS	86480
LABCORP	016774	Q Fever Antibodies, IgG Container:Red-top tube or gel-barrier tube Temperature:Maintain specimen at room temperature.	86638
LABCORP	806606	Q Fever IgG/IgM Antibody Scrn Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x4
LABCORP	806617	Q Fever IgM Antibody Screen Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x2
LABCORP	199995	QAPAM	NOCPT
LABCORP	182913	QuantiFERON-TB Gold Plus Container:Green-Top (Lithium Heparin) Tube Temperature:After collection, the blood can be held at room temperature (17 to 25 degrees Celsius) for up to three hours, and then placed at 2 to 8 degrees Celsius for up to 48 hours. Alternatively, blood can immediately be placed at 2 to 8 degrees Celsius for up to 48 hours. *After refrigeration, the lithium heparin tube MUST equilibrate to room temperature (17 to 25 degrees Celsius) prior to the transfer to QFT-Plus Blood Collection Tubes. *Aliquoted QFT-Plus Blood Collection Tubes should be placed in the 37 degree Celsius incubator within two hours of blood transfer. *NOTE: Total time from blood draw to incubation in QFT-Plus Blood Collection Tubes should NOT exceed 53 hours.	86480
LABCORP	808404	Quetiapine, Serum or Plasma Container:Red-top tube OR green-top (heparin) tube. Gel-barrier tubes are NOT recommended. Temperature:SUBMISSION/TRANSPORT (<3 days): Room Temperature. For storage beyond three days, specimen should be refrigerated or frozen.	80299
LABCORP	791462	Quetiapine, Ur Container:Urine transfer tube Temperature:Maintain specimen at room temperature up to three days. Specimens may also be refrigerated or frozen.	80342
LABCORP	007831	Quinidine, Serum Container:Red-top tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant. Temperature:Room Temperature	80194
LABCORP	810819	Quinine, Serum/Plasma Container:Red-top tube or lavender-top (EDTA) tube Temperature:Refrigerate	84228
LABCORP	802110	Quinine/Quinidine Different. Container:Red-top tube, lavender-top (EDTA) tube Temperature:Refrigerate (Cool Pack)	80194
LABCORP	805913	Quinoa (Chenopodium quinoa)IgE Container:Gel-barrier tube Temperature:Maintain specimen at room temperature	86003
LABCORP	827488	Quinoline Yellow - BasoFu. HRT Container:Green-top (sodium heparin) tube: submit original tube Temperature:Maintain specimen at room temperature	86343