

# PMG Laboratory Spokane

## Test Directory

| LAB     | CODE    | TEST NAME   | CPT4 CODE |
|---------|---------|---|-----------|
| PMGLAB  | LAB560  | QUAD SCREEN MATERNAL, SERUM<br>Place Of Service:LABCORP RTP   |           |
| PMGLAB  | LAB2399 | QUANTIFERON GOLD<br>Container:Dark green lithium heparin tube<br>Temperature:Refrigerate<br>Place Of Service:PMG-PCS  | 86480     |
| LABCORP | 016774  | Q Fever Antibodies, IgG<br>Container:Red-top tube or gel-barrier tube<br>Temperature:Maintain specimen at room temperature.   | 86638     |
| LABCORP | 806606  | Q Fever IgG/IgM Antibody Scrn<br>Container:Red-top tube or gel-barrier tube<br>Temperature:Refrigerate  | 86638 x4  |
| LABCORP | 806617  | Q Fever IgM Antibody Screen<br>Container:Red-top tube or gel-barrier tube<br>Temperature:Refrigerate  | 86638 x2  |
| LABCORP | 199995  | QAPAM   | NOCPT     |
| LABCORP | 182913  | QuantiFERON-TB Gold Plus<br>Container:Green-Top (Lithium Heparin) Tube<br>Temperature:After collection, the blood can be held at room temperature (17 to 25 degrees Celsius) for up to three hours, and then placed at 2 to 8 degrees Celsius for up to 48 hours. Alternatively, blood can immediately be placed at 2 to 8 degrees Celsius for up to 48 hours. *After refrigeration, the lithium heparin tube MUST equilibrate to room temperature (17 to 25 degrees Celsius) prior to the transfer to QFT-Plus Blood Collection Tubes. *Aliquoted QFT-Plus Blood Collection Tubes should be placed in the 37 degree Celsius incubator within two hours of blood transfer. *NOTE: Total time from blood draw to incubation in QFT-Plus Blood Collection Tubes should NOT exceed 53 hours. | 86480     |
| LABCORP | 808404  | Quetiapine, Serum or Plasma<br>Container:Red-top tube OR green-top (heparin) tube. Gel-barrier tubes are NOT recommended.<br>Temperature:SUBMISSION/TRANSPORT (<3 days): Room Temperature. For storage beyond three days, specimen should be refrigerated or frozen.  | 80299     |
| LABCORP | 791462  | Quetiapine, Ur<br>Container:Urine transfer tube<br>Temperature:Maintain specimen at room temperature up to three days. Specimens may also be refrigerated or frozen.  | 80342     |
| LABCORP | 007831  | Quinidine, Serum<br>Container:Red-top tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant.<br>Temperature:Room Temperature  | 80194     |
| LABCORP | 810819  | Quinine, Serum/Plasma<br>Container:Red-top tube or lavender-top (EDTA) tube<br>Temperature:Refrigerate  | 84228     |
| LABCORP | 802110  | Quinine/Quinidine Different.<br>Container:Red-top tube, lavender-top (EDTA) tube<br>Temperature:Refrigerate (Cool Pack)   | 80194     |
| LABCORP | 805913  | Quinoa (Chenopodium quinoa)IgE<br>Container:Gel-barrier tube<br>Temperature:Maintain specimen at room temperature   | 86003     |
| LABCORP | 827488  | Quinoline Yellow - BasoFu. HRT<br>Container:Green-top (sodium heparin) tube: submit original tube<br>Temperature:Maintain specimen at room temperature  | 86343     |