

Online Test Menu Laboratory Catalog



LAB	CODE	TEST NAME	CPT4 CODE
SPH	LAB1209	Q FEVER AB Place Of Service:LabCorp	
SPH	LAB560	QUAD SCREEN MATERNAL, SERUM Place Of Service:LabCorp	
SPH	LAB2399	Quantiferon TB Gold Plus Place Of Service:LabCorp	
SPH	LAB23381	QUETIAPINE, UR Place Of Service:MEDTOX LABORATORIES INC	
LABCORP	806606	Q Fever IgG/IgM Antibody Scrn Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x4
LABCORP	806617	Q Fever IgM Antibody Screen Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x2
LABCORP	182893	QuantiFERON(R)-TB Gold Plus (Client Incubated) Container:The QuantiFERON(R)-TB Gold Plus collection kit contains the instructions for the collection and handling of (one each): (1) gray-top (with white ring), uncoated (nil); (2) green cap with white ring, TB1 tube; (3) yellow cap with white ring, TB2 tube; (4) purple top with white ring, mitogen-coated. A high altitude kit is also available for locations between 3350 and 6150 feet; a cap with a yellow ring differentiates the tubes. Temperature:1. Maintain specimen at room temperature prior to incubation at 37 degrees C. Stability: Kit MUST be incubated within 16 hours of collection. 2. After incubation, maintain specimen at room temperature (17 degrees C to 27 degrees C). Do NOT centrifuge refrigerate, or ship tubes on ice.	86480
LABCORP	182913	QuantiFERON-TB Gold Plus Container:Green-Top (Lithium Heparin) Tube Temperature:After collection, the blood can be held at room temperature (17 to 25 degrees Celsius) for up to three hours, and then placed at 2 to 8 degrees Celsius for up to 48 hours. Alternatively, blood can immediately be placed at 2 to 8 degrees Celsius for up to 48 hours. *After refrigeration, the lithium heparin tube MUST equilibrate to room temperature (17 to 25 degrees Celsius) prior to the transfer to QFT-Plus Blood Collection Tubes. *Aliquoted QFT-Plus Blood Collection Tubes should be placed in the 37 degree Celsius incubator within two hours of blood transfer. *NOTE: Total time from blood draw to incubation in QFT-Plus Blood Collection Tubes should NOT exceed 53 hours.	86480