

Online Test Menu

LABORATORY CATALOG

LAB	CODE	TEST NAME	CPT4 CODE
BSHSI-R	QFIL	QUANTIFERON-TB PLUS(CLIENT INCUB.)	
LABCORP	806606	Q Fever IgG/IgM Antibody Scrn Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x4
LABCORP	806617	Q Fever IgM Antibody Screen Container:Red-top tube or gel-barrier tube Temperature:Refrigerate	86638 x2
LABCORP	199995	QAPAM	NOCPT
LABCORP	182879	QuantiFERON(R)-TB Gold Plus Container:The QuantiFERON(R)-TB Gold Plus collection kit contains the instructions for the collection and handling of (one each): (1) gray-top (with white ring), uncoated (nil); (2) green cap with white ring, TB1 tube; (3) yellow cap with white ring, TB2 tube; (4) purple top with white ring, mitogen-coated. A high altitude kit is also available for locations between 3350 and 6150 feet; a cap with a yellow ring differentiates the tubes. Temperature:Maintain specimen at room temperature for four tube collection kit. Stability: Kit MUST be incubated within 16 hours of collection.	86480
LABCORP	182913	QuantiFERON-TB Gold Plus Container:Green-Top (Lithium Heparin) Tube Temperature:After collection, the blood can be held at room temperature (17 to 25 degrees Celsius) for up to three hours, and then placed at 2 to 8 degrees Celsius for up to 48 hours. Alternatively, blood can immediately be placed at 2 to 8 degrees Celsius for up to 48 hours. *After refrigeration, the lithium heparin tube MUST equilibrate to room temperature (17 to 25 degrees Celsius) prior to the transfer to QFT-Plus Blood Collection Tubes. *Aliquoted QFT-Plus Blood Collection Tubes should be placed in the 37 degree Celsius incubator within two hours of blood transfer. *NOTE: Total time from blood draw to incubation in QFT-Plus Blood Collection Tubes should NOT exceed 53 hours.	86480
LABCORP	791462	Quetiapine, Ur Container:Urine transfer tube Temperature:Maintain specimen at room temperature up to three days. Specimens may also be refrigerated or frozen.	80342
LABCORP	007831	Quinidine, Serum Container:Red-top tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant. Temperature:Room Temperature	80194
LABCORP	810819	Quinine, Serum/Plasma Container:Red-top tube or lavender-top (EDTA) tube Temperature:Refrigerate	84228
LABCORP	802110	Quinine/Quinidine Different. Container:Red-top tube, lavender-top (EDTA) tube Temperature:Refrigerate (Cool Pack)	80194
LABCORP	805913	Quinoa (Chenopodium quinoa)IgE Container:Gel-barrier tube Temperature:Maintain specimen at room temperature	86003
LABCORP	827488	Quinoline Yellow - BasoFu. HRT Container:Green-top (sodium heparin) tube: submit original tube Temperature:Maintain specimen at room temperature	86343